

EUCALYPT

VEGETATION TYPE 1


Regional Ecosystem: 12.11.24

Broad-leaved White Mahogany - Queensland Stringybark (*Eucalyptus carnea* - *E. tindaliae*) Woodland to Open Forest on Metasediments



COMMUNITY STRUCTURE

Vegetation type (VT) 1 is typically a woodland to open forest providing approximately 25-50% canopy cover (% shade to underlying plants). The canopy layer is typically 15-27m high, with *Eucalyptus carnea* and *E. tindaliae* the most regularly occurring tree types, although a wide variety of other species (most notably *Corymbia intermedia*, *C. citriodora*, *C. henryi*, *E. siderophloia*, and *E. carnea*) can be present or locally dominant.



The sub-canopy and shrub layers below the canopy are usually sparse (ca. 0-10% cover) with wattles (*Acacia disparrima*, *A. concurrens*), she-oaks (*Allocasuarina torulosa*, *C. littoralis*) and eucalypt saplings the most commonly occurring low trees and shrubs. The ground layer is dominated by grasses (particularly *Themeda triandra*, *Imperata cylindrica* and *Cymbopogon refractus*, although *Lepidosperma laterale* and *Lomandra laxa* are occasionally dominant).

Characteristic plant species

Approximately **127 native plants** species have been recorded for this vegetation type. Characteristic plant species for this vegetation type are listed below. Dominant (most numerous) species are shaded.



Indicates species is a preferred koala food tree*



Indicates species is a Glossy Black-Cockatoo feed tree species



Indicates species is a City-wide significant species

* It is noted that in addition to preferred food trees, koalas utilise a range of eucalypt and non-eucalypt tree species for supplemental feeding and other uses such as shelter. These other species are also important and necessary features of koala habitat.

CANOPY

Upper layer of vegetation exposed to sunlight which creates a canopy that shades lower layers




Lui Weber ©

Broad-leaved White Mahogany
Eucalyptus carnea



Queensland White Stringybark
Eucalyptus tindaliae



Pink Bloodwood
Corymbia intermedia



Grey Ironbark
Eucalyptus siderophloia



Terra Ark ©

CANOPY (CON'T)

Upper layer of vegetation exposed to sunlight which creates a canopy that shades lower layers



Small-fruited Grey Gum
Eucalyptus propinqua



City of Gold Coast ©



Spotted Gum
Corymbia citriodora subsp. *variegata*




Photo needed



Narrow-leaved Ironbark
Eucalyptus crebra



Smooth-barked Apple
Angophora leiocarpa



Tallowood
Eucalyptus microcorys



Forest Red Gum
Eucalyptus tereticornis



Narrow-leaved Red Gum
Eucalyptus seeana



Large-leaved Spotted Gum
Corymbia henryi

SUB-CANOPY

Tree layer below canopy. Small trees and juvenile canopy trees.



Brush Box
Lophostemon confertus



Forest She-Oak
Allocasuarina torulosa



Black She-oak
Allocasuarina littoralis



Hickory Wattle
Acacia disparrima subsp.
disparrima



Black Wattle
Acacia concurrens



Swamp Box
Lophostemon suaveolens



Pink Bloodwood
Corymbia intermedia

SHRUB LAYER

Middle layer of vegetation usually made up of small trees (including juvenile canopy and sub canopy tree species) and woody shrubs



Juvenile tree

Hickory Wattle

Acacia disparrima subsp. disparrima



Juvenile tree

Black Wattle

Acacia concurrens



GBC

Juvenile tree

Forest She-Oak

Allocasuarina torulosa



Juvenile tree

Brush Box

Lophostemon confertus



GBC

Juvenile tree

Black She-oak

Allocasuarina littoralis



Brisbane Wattle/Fringed Wattle

Acacia fimbriata



Dogwood

Jacksonia scoparia



Juvenile tree

Swamp Box

Lophostemon suaveolens

GROUND LAYER

Lowest layer of vegetation. Plant types can include grasses; graminoids (non-woody plants with a grass-like morphology); ferns; and forbs (non-woody, broad-leaved, flowering plants). Vines which extend upwards into the canopy are uncommon.



Kangaroo Grass
Themeda triandra
GRASS (TUSSOCK)




Blady Grass
Imperata cylindrica
GRASS




Barbwire Grass
Cymbopogon refractus
GRASS



Poverty Grass
Eremochloa bimaclulata
GRASS (TUSSOCK)



Small-flowered Fingergrass
Digitaria parviflora
GRASS (TUSSOCK)



Wiry Panic
Entolasia stricta
GRASS (TUSSOCK)



Graceful / Pademelon Grass
Ottochloa gracillima
GRASS



Weeping Grass
Microlaena stipoides
GRASS (TUSSOCK)



Variable Sword-sedge
Lepidosperma laterale
GRAMINOID (SEGE)



Broad-leaved Mat Rush
Lomandra laxa
GRAMINOID

Graham McDonald ©

City-wide significant plant species



The City of Gold Coast recognises species which are locally significant as City-wide significant (CWS) species. These species are important because they may be threatened, restricted to the Gold Coast, or at the edge of their geographic range. In addition to characteristic species identified above as CWS species, the following CWS plant species may also be present in this vegetation type.



Rainforest Carex
Carex breviculmis
SEDGE



Dysentery Plant
Grewia latifolia
SHRUB




Cabbage Tree Palm
Livistona australis
PALM

OCCURRENCE

Native plants occur in vegetation communities, which are consistently associated with a particular soil type, landform (shape of the land, e.g. hills or plains) aspect (position on a slope in relation to the sun) and climate.

This vegetation type occurs on lower hillslopes and gently undulating country at the base of slopes through the northern to central parts of the Gold Coast. It also occurs on higher footslopes in more fertile areas around Mudgeeraba in the south of the City. Generally occurring on thin to mid-depth soils with some litter and humus development, on mudstone and shale-based soils on lower foothill slopes, where it transitions into VT4 (Spotted Gum/Ironbark Woodland) on higher slopes and thinner more gravelly soils. Common localities include Coomera, Nerang, Maudsland, Pimpama, Ormeau and Mudgeeraba.

Historic distribution of Vegetation Type 1



* Metasediment rocks

The most common underlying geology on the Gold Coast is metasediment rocks. Metasediment rocks are a type of metamorphic rock (rock transformed by heat and pressure). Originally these rocks were sedimentary rocks which were formed on the ocean floor through the deposition and solidification of sediment. These sedimentary rocks were subsequently buried underneath other rocks and subjected to high pressures and temperatures, causing the rock to recrystallize. This recrystallization process is known as metamorphism, hence the term metamorphic rocks. About 300 million years ago these metamorphic rocks were pushed upward by geologic processes, creating much of the ranges, hills and lowlands on the Gold Coast.

2017 EXTENT AND CONSERVATION STATUS

Gold Coast


Historically, this vegetation type was the most common vegetation type on the Gold Coast and remains so today, with 52% of its original extent remaining. The 2017 extent* was 11,281 hectares.

1 HECTARE (HA) = 2.46 ACRES = THE SIZE OF AN INTERNATIONAL RUGBY FIELD

EXTENT (ha)

Historic
21,895ha

2015*
11,281ha
52% of
historical
extent




* Extent as mapped in 2017. Includes remnant vegetation only. Does not include disturbed remnant or regrowth

Queensland

The conservation status of vegetation in Queensland is specified under the Vegetation Management Act 1999, which lists this regional ecosystem (RE 12.11.24) as being 'Least Concern'.

LIKELIHOOD OF BECOMING EXTINCT (in QLD) due to biodiversity loss/degradation



USEFUL RESOURCES

City of Gold Coast website: Environmental weeds and invasive plants.

Find out more about regional ecosystems at the Queensland Government Regional Ecosystems webpage.

CREDITS

Content – ngh Environmental and Jason Searle.
Vegetation Type Photo – Lui Weber ©
Unless otherwise noted all other photos – Glenn Leiper ©

Version 3, November 2020

THREATS

This vegetation type is generally quite open. The ground layer is typically a mosaic of grassland and more open areas with a conspicuous litter layer of leaves and bark. Too frequent fires, over-grazing or other loss of native grassy understorey can result in erosion and desiccation of the soil layer. Fire management, particularly where the more common weeds of Lantana and Molasses Grass have become established, is important in weed control and in managing the soils of these areas. Please refer to the Queensland Government's Regional Ecosystem Description Database for more information including suggested fire regimes.

Common threats to all vegetation types

Clearing

Native vegetation is protected by Federal, State and local legislation. However, with increasing population growth in the region, Southeast Queensland is experiencing large amounts of vegetation clearing, particularly in areas designated for urban development. Protecting native vegetation on your property is one of the most beneficial things you can do to protect wildlife and the natural environment.

Weeds

Environmental weeds are the second biggest threat to our natural environment after land clearing. Environmental weeds (introduced plants that have naturalised and are invading our bushland) degrade our natural environment by:

- out competing native plant species for available nutrients and light,
- taking over and transforming native landscapes often leading to local plant or animal extinctions and loss of biodiversity,
- reducing the availability of food and other resources for many native animals whilst sometimes benefiting pest animals,
- increasing the risk of destructive wildfire, and/or
- often being toxic to people and animals.

Fire

Very broadly, vegetation types are either adapted to fire or fire sensitive. Fire can become a threat if:

- it extends into vegetation types which should not be burnt e.g. rainforest,
- the frequency and/or intensity of the fire is too high,
- the area burnt is too large.

Grazing

The grazing of animals like cattle, horses, goats and feral animals such as deer can cause trampling or loss of diversity of seedlings and compact soil, preventing natural regeneration.

Collecting

Unethical and illegal collection of plant specimens in the wild poses a serious threat to some species, particularly orchids, grass trees and epiphytes.

Climate change

Changes in temperature and rainfall can have significant effects on our city's vegetation. For example, without consistent rainfall, areas become drier, potentially resulting in higher fire frequency and/or intensity, which some plants and vegetation communities won't be able to tolerate. Plants (and animals) need available space to migrate as conditions change, with high altitude species at the greatest risk as there is nowhere suitable for them to go. Warmer conditions may also provide the right habitat for a greater variety of weeds. As sea levels rise, salt water moves further upstream and vegetation also becomes inundated.