

EUCALYPT

VEGETATION TYPE 1a


Regional Ecosystem: 12.11.3

White Mahogany +/- Tallowwood - Grey Gum
(*Eucalyptus acmenoides* +/- *E. microcorys* - *E. propinqua*) Open Forest on Metasediments



COMMUNITY STRUCTURE

Vegetation type (VT) 1a is typically a tall open forest adjoining rainforest. It has a relatively dense canopy (60-65% cover*). The canopy layer is typically 19-32m high, with White Mahogany (*Eucalyptus acmenoides*) often dominant, with *E. microcorys*, *E. propinqua* and *E. siderophloia* also often present.



The sub-canopy is sparse and mainly composed of Brush Box (*Lophostemon confertus*) and eucalypt saplings. The shrub layer is sparse yet diverse in some locations. The ground layer is dominated by grasses and/or ferns (particularly *Entolasia stricta*, *Cymbopogon refractus*, *Imperata cylindrica*, *Doodia aspera* and *Blechnum cartilagineum*), with a diverse array of sub-shrubs (including *Solanum stelligerum* and *Zieria collina*) also present. Vines are uncommon. Where fire is excluded, ferns, cycads and other rainforest plants like gingers become more prolific

*canopy covers is % shade to underlying plants

Characteristic plant species

Approximately **84 native plants** species have been recorded for this vegetation type. Characteristic plant species for this vegetation type are listed below. Dominant (most numerous) species are shaded.



Indicates species is a preferred koala food tree*



Indicates species is a Glossy Black-Cockatoo feed tree species



Indicates species is a City-wide significant species

* It is noted that in addition to preferred food trees, koalas utilise a range of eucalypt and non-eucalypt tree species for supplemental feeding and other uses such as shelter. These other species are also important and necessary features of koala habitat.

CANOPY

Upper layer of vegetation exposed to sunlight which creates a canopy that shades lower layers



White Mahogany
Eucalyptus acmenoides



Tallowwood
Eucalyptus microcorys



Small-fruited Grey Gum
Eucalyptus propinqua



Grey Ironbark
Eucalyptus siderophloia



Pink Bloodwood
Corymbia intermedia



Brush Box
Lophostemon confertus



SUB-CANOPY

Tree layer below canopy



Forest She-Oak
Allocasuarina torulosa



Brush Box
Lophostemon confertus




Hickory Wattle
Acacia disparrima subsp. disparrima

SHRUB LAYER

Middle layer of vegetation usually made up of small trees (including juvenile canopy and sub canopy tree species) and woody shrubs



Hickory Wattle
Acacia disparrima subsp. disparrima



Devils Needles
Solanum stelligerum



Mt Tamborine Ziera *Zieria collina*



Golden Pea
Daviesia arborea



Waddy Wood *Trochocarpa laurina*

GROUND LAYER AND VINES

Lowest layer of vegetation. Plant types can include grasses; graminoids (non-woody plants with a grass-like morphology); ferns; and forbs (non-woody, broad-leaved, flowering plants). Grasses constitute most of the ground cover although ferns are prominent in some cases. Vines are uncommon.



Gristle Fern
Blechnum cartilagineum

FERN



Kangaroo Grass
Themeda triandra

TUSOCK GRASS



Blady Grass
Imperata cylindrica

GRASS



Bluegrass
Poa cheelii

GRASS



Wiry Panic
Entolasia stricta

TUSOCK GRASS



Barbwire Grass
Cymbopogon refractus

GRASS








Prickly Rasp Fern
Doodia aspera

FERN

City-wide significant plant species



The City of Gold Coast recognises species which are locally significant as City-wide significant (CWS) species. These species are important because they may be threatened, restricted to the Gold Coast, or at the edge of their geographic range. In addition to characteristic species identified above as CWS species, the following CWS plant species may also be present in this vegetation type.


		
<p>Ironbark Orchid <i>Dendrobium aemulum</i> EPIPHYTE</p>	<p>Rainforest Carex <i>Carex breviculmis</i> GRASS</p>	<p>Palm Lily <i>Cordyline congesta</i> PALM</p>
		
<p>Shirley's Nightshade <i>Solanum shirleyanum</i> SHRUB</p>	<p>Tongue Orchid <i>Dockrillia linguiformis</i> EPIPHYTE</p>	

OCCURRENCE

Native plants occur in vegetation communities, which are consistently associated with a particular soil type, landform (shape of the land, e.g. hills or plains) aspect (position on a slope in relation to the sun) and climate.

This vegetation type occurs on higher hillslopes and escarpment areas on deep fertile soils, typically with deep leaf litter, high organic and good moisture content. This community is largely restricted to the Wongawallan and Mt Tamborine areas in the central western parts of Gold Coast City, often adjoining rainforest.

Historic distribution of Vegetation Type 1a



2017 EXTENT AND CONSERVATION STATUS

Gold Coast

The 2017 extent* of this vegetation type on the Gold Coast is 2,524 hectares


1 HECTARE (HA) = 2.46 ACRES = THE SIZE OF AN INTERNATIONAL RUGBY FIELD

EXTENT (ha)

Historic
3,210ha

2017*
2,523ha

79% of
historical
extent




* Extent as mapped in 2017. Includes remnant vegetation only. Does not include disturbed remnant or regrowth

Queensland

The conservation status of vegetation in Queensland is specified under the Vegetation Management Act 1999, which lists this regional ecosystem (RE 12.11.3) as being of 'Least Concern'.

LIKELIHOOD OF BECOMING EXTINCT (in QLD) due to biodiversity loss/degradation



USEFUL RESOURCES

City of Gold Coast website: Environmental weeds and invasive plants.

Find out more about regional ecosystems at the Queensland Government Regional Ecosystems webpage.

CREDITS

Content – ngh Environmental and Jason Searle.
Vegetation Type Photo – Lui Weber ©
Unless otherwise noted all other photos – Glenn Leiper ©

Version 3, November 2020

THREATS

This vegetation type is largely restricted to more fertile escarpment areas adjoining rainforest and is susceptible to infestation by Lantana and other dense understorey growth, particularly in the absence of fire. It is a naturally restricted vegetation type that can transition towards rainforest in the absence of fire, and relies on appropriate fire management (low frequency, moderate to high intensity fires). Weed and fire management are key to the healthy management of this vegetation type. Please refer to the Queensland Government's Regional Ecosystem Description Database for more information including suggested fire regimes.

Common threats to all vegetation types

Clearing

Native vegetation is protected by Federal, State and local legislation. However, with increasing population growth in the region, Southeast Queensland is experiencing large amounts of vegetation clearing, particularly in areas designated for urban development. Protecting native vegetation on your property is one of the most beneficial things you can do to protect wildlife and the natural environment.

Weeds

Environmental weeds are the second biggest threat to our natural environment after land clearing. Environmental weeds (introduced plants that have naturalised and are invading our bushland) degrade our natural environment by:

- out competing native plant species for available nutrients and light,
- taking over and transforming native landscapes often leading to local plant or animal extinctions and loss of biodiversity,
- reducing the availability of food and other resources for many native animals whilst sometimes benefiting pest animals,
- increasing the risk of destructive wildfire,
- often being toxic to people and animals.

Fire

Very broadly, vegetation types are either adapted to fire or fire sensitive. Fire can become a threat if:

- it extends into vegetation types which should not be burnt e.g. rainforest,
- the frequency and/or intensity of the fire is too high,
- the area burnt is too large.

Grazing

The grazing of animals like cattle, horses, goats and feral animals such as deer can cause trampling or loss of diversity of seedlings and compact soil, preventing natural regeneration.

Collecting

Unethical and illegal collection of plant specimens in the wild poses a serious threat to some species, particularly orchids, grass trees and epiphytes.

Climate change

Changes in temperature and rainfall can have significant effects on our city's vegetation. For example, without consistent rainfall, areas become drier, potentially resulting in higher fire frequency and/or intensity, which some plants and vegetation communities won't be able to tolerate. Plants (and animals) need available space to migrate as conditions change, with high altitude species at the greatest risk as there is nowhere suitable for them to go. Warmer conditions may also provide the right habitat for a greater variety of weeds. As sea levels rise, salt water moves further upstream and vegetation also becomes inundated.